



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/10, C12P 19/34	A1	(11) International Publication Number: WO 96/36706 (43) International Publication Date: 21 November 1996 (21.11.96)
(21) International Application Number: PCT/US96/07083 (22) International Filing Date: 15 May 1996 (15.05.96) (30) Priority Data: 08/446,118 19 May 1995 (19.05.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/446,118 (CIP) Filed on 19 May 1996 (19.05.96) (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): LEE, Ann, L. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). SAGAR, Sangeetha [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		(81) Designated States: AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IS, JP, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: A METHOD FOR LARGE SCALE PLASMID PURIFICATION (57) Abstract <p>A process is disclosed for the large scale isolation and purification of plasmid DNA from large scale microbial fermentations. All three forms of plasmid DNA; supercoil (form I), nicked or relaxed circle (form II), and linearized (form III), are individually isolatable using the disclosed process. Highly purified DNA suitable for inclusion in a pharmaceutical composition is provided by the disclosed process.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

TITLE OF THE INVENTION

A METHOD FOR LARGE SCALE PLASMID PURIFICATION

RELATED APPLICATION

5 This is a continuation-in-part application of U.S. Serial
Number 08/446,118 filed May 19, 1995.

BACKGROUND OF THE INVENTION

10 The classical techniques for isolating plasmid DNA from
microbial fermentations are suitable for small or laboratory scale
plasmid preparations. One such procedure involves the alkaline lysis of
microbial host cells containing the plasmid, followed by acetate
neutralization causing the precipitation of host cell genomic DNA and
15 proteins which are then removed by, for example, centrifugation. The
liquid phase contains the plasmid DNA which is alcohol precipitated and
then subjected to isopycnic centrifugation using CsCl in the presence of
ethidium bromide. The ethidium bromide is required in order to
20 separate the total plasmid DNA into the three different forms,
supercoiled (form I), nicked circle (form II), and linearized (form III),
and the desired plasmid form is collected. Further extraction with
butanol is required to remove residual ethidium bromide followed by
DNA precipitation using alcohol. Additional purification steps follow
25 to remove host cell proteins. The removal of host proteins is
performed by repeated extractions using phenol or a mixture of phenol
and chloroform. The plasmid DNA is alcohol precipitated and residual
phenol is removed by repeated isoamyl/chloroform extractions. The
final alcohol precipitated plasmid DNA is dissolved in water or a
suitable buffer solution.

30 There are numerous drawbacks and limitations to this
process including:

- a) this process requires the use of expensive and hazardous
chemicals (CsCl and EtBr, which are used within the
density gradient centrifugation; EtBr is a known mutagen

- 2 -

and must be removed from products; also it is an intercalating agent which can nick the plasmid);

- 5 b) the density centrifugation step is not easily scaleable;
- c) there is a need for organic solvent extraction to remove residual EtBr;
- d) phenol extraction is used to remove residual proteins and DNase, a process that would require a centrifuge to break
10 phenol/water emulsion;
- e) highly repetitive steps making it laborious and time consuming (isolation requires several days);
- 15 f) scalability of the chemical lysis step is an obstacle i.e., lysozyme/alkaline/KOAc treatment step is efficient in lysing cells on a small scale, however, the increase in viscosity makes large scale processing very difficult; and
- 20 g) use of large quantities of lysozyme to enzymatically weaken the microbial cell wall prior to lysis.

The mixture is then neutralized by addition of acid which results in precipitation of the high molecular weight chromosomal
25 DNA. The high molecular weight RNA and protein-SDS complexes precipitate with the addition of high concentration of KOAc salt. The plasmid product remains in the clarified supernatant following centrifugation. Limitations here include the need to process quickly and on ice in order to retard the activity of nucleases which are not removed
30 until phenol extraction. The main contaminant remaining in the supernatant with the product is RNA.

Another commonly utilized method for isolating and purifying plasmid DNA from bacteria provides a rapid process suitable for only very small scale preparations.

- 3 -

Holmes and Quigley (1981, *Analytical Biochem.*, **114**, pp 193-197) reported a simple and rapid method for preparing plasmids where the bacteria are treated with lysozyme, then boiled at about 100°C in an appropriate buffer (STET) for 20 - 40 seconds forming an insoluble clot of genomic DNA, protein and debris leaving the plasmid in solution with RNA as the main contaminant. Lysozyme is apparently a requirement for this technique to work, and as such adds a treatment step which is less desirable for large scale manufacture of DNA for human or veterinary use. However, the addition of lysozyme may enhance plasmid release during lysis. An advantage is that heat treatment of the cells would also denature the DNase. However, this technique is not suitable for scale up to a high volume of microbial fermentations and is meant for fermentations less than five liters.

Alternatives to isopycnic centrifugation using CsCl for plasmid purification have been published. These alternatives are suitable only for laboratory scale plasmid isolation and include:

- a) size exclusion chromatography, which is inherently limited in throughput;
- b) hydroxyapatite chromatography, which has the disadvantage of requiring high concentrations of urea for efficiency;
- c) reversed phase chromatography; and
- d) ion exchange chromatography.

Large scale isolation and purification of plasmid DNA from large volume microbial fermentations, therefore, requires the development of an improved plasmid preparation process. An isolation and purification process for large scale plasmid DNA production is necessitated by recent developments in many areas of molecular biology. In particular, recent advances in the field of polynucleotide-

- 4 -

based vaccines for human use, and potentially human gene therapy, requires the ability to produce large quantities of the polynucleotide vaccine in a highly purified form.

- Unprecedented technology is required for developing/-
5 implementing a large scale commercially viable process for fermentation, isolation, purification and characterization of DNA as a biopharmaceutical.

SUMMARY OF THE INVENTION

- 10 The current laboratory method used to isolate and purify plasmid DNA consists of a series of classical laboratory techniques that are not suitable for a manufacturing process. For example, density gradient centrifugations are not scaleable; the purification procedure necessitates the use of hazardous and expensive chemicals/solvents such
15 as ethyidium bromide, a known mutagen, and is labor intensive and time consuming. Therefore, a scaleable alternative process was developed, and is disclosed herein. In addition, an HPLC assay was established to track the plasmid product through the process steps and to distinguish between the plasmid forms. The microbial cells harboring the plasmid
20 are suspended and optionally incubated with lysozyme in a buffer containing detergent, heated using a flow-through heat exchanger to lyse the cells, followed by centrifugation. After centrifugation the clarified lysate, which contains predominately RNA and the plasmid product, is filtered through a 0.45 micron filter and then diafiltered, prior to
25 loading on the anion exchange column. The plasmid product may optionally be treated with RNase before or after filtration, or at an earlier or later step. The anion exchange product fraction containing the plasmid is loaded onto the reversed phase column, and is eluted with an appropriate buffer, providing highly pure plasmid DNA suitable for
30 human use.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. A schematic of a suitable heat exchanger apparatus is shown.

- 5 -

- Figure 2. The relationship between the outlet temperature and the flow rate is shown, graphically.
- 5 Figure 3. Comparative chromatograms of total plasmid in clarified supernatant with 50 mM EDTA and 100 mM EDTA are shown.
- 10 Figure 4. The yield of supercoiled plasmid as a function of exit temperature is shown.
- Figure 5. The elution profiles of anion exchange columns run with RNase treated (bold line) and untreated (thin line) clarified lysates are shown.
- 15 Figure 6. The elution profiles of anion exchange chromatography with clarified lysate that was diafiltered before the column or not diafiltered before the column are shown.
- 20 Figure 7. An elution profile of plasmid DNA from cell lysate is shown.
- Figure 8. An agarose gel electrophoresis analysis of the DNA product obtained at various intermediate steps of purification is shown.
- 25 Figure 9. A tracing of the anion exchange HPLC analysis of the DNA product demonstrating the purity of the product is shown.

DETAILED DESCRIPTION OF THE INVENTION

- 30 We have identified a novel, scaleable, alternative lysis/debris removal process for large scale plasmid isolation and purification which exploits a rapid heating method to induce cell lysis and precipitate genomic DNA, proteins and other debris while keeping the plasmid in solution. The utility of this process is the large scale

- 6 -

isolation and purification of plasmid DNA. We have found that suspending the microbial cells in modified STET buffer (described below) and then heating the suspension to about 70-100°C in a flow-through heat exchanger results in excellent lysis. Continuous flow or
5 batch-wise centrifugation of the lysate effects a pellet that contains the cell debris, protein and most of the genomic DNA while the plasmid remains in the supernatant. This invention offers a number of advantages including higher product recovery than by chemical lyses, inactivation of DNases, operational simplicity and scalability.

10 The present invention is drawn to a process for the large scale isolation and purification of plasmid DNA from microbial fermentations. Large scale microbial cell fermentations as used herein are considered to be total cell fermentation volumes of greater than about 5 liters, or the cells harvested from a fermentation volume
15 greater than about 5 liters.

The present invention is also drawn to providing plasmid DNA in a highly purified form suitable for human use. DNA for human use includes, but is not limited to, polynucleotide vaccines and DNA for human gene therapy. Polynucleotide vaccines are intended for
20 direct injections into humans [Montgomery, D.L. *et al.*, 1993, *Cell Biol.*, 169, pp. 244-247; Ulmer, J.B. *et al.*, 1993, *Science*, 259, pp. 1745-1749].

The present invention is also drawn to an in-line monitoring process for the tracking of the various forms of plasmid
25 DNA through the isolation and purification steps. The various forms of plasmid DNA referred to above which can be individually isolated by the process of the present invention are form I (supercoiled plasmid), form II (nicked or relaxed plasmid), and form III (linearized plasmid).

The process of the present invention is suitable for use with
30 microbial fermentations in general. It is readily apparent to those skilled in the art that a wide variety of microbial cells are suitable for use in the process of the present invention, including but not limited to, fungal cells including yeast, and bacterial cells. A preferred microbial fermentation is a bacterial fermentation of cells containing the plasmid

- 7 -

to be isolated and purified. A preferred bacterial fermentation is a fermentation of *E. coli* containing the plasmid to be isolated and purified. It is readily apparent to those skilled in the art that bacterial fermentations other than *E. coli* fermentations are suitable for use in the present invention. The microbial fermentation may be grown in any liquid medium which is suitable for growth of the bacteria being utilized.

The plasmid to be isolated and purified by the process of the present invention can be any extrachromosomal DNA molecule. The plasmids can be high copy number per cell or low copy number per cell. The plasmids can also be of virtually any size. It is readily apparent to those skilled in the art that virtually any plasmid in the microbial cells can be isolated by the process of the present invention.

Microbial cells containing the plasmid are harvested from the fermentation medium to provide a cell paste, or slurry. Any conventional means to harvest cells from a liquid medium is suitable, including, but not limited to centrifugation or microfiltration.

Isolation of the plasmid DNA from harvested microbial cells using the current lab scale procedures consist mainly of enzymatic treatment of microbial cells to weaken the cell wall followed by cell lysis. The purification steps include repetitive CsCl/EtBr centrifugations followed by organic solvent extractions and precipitation to remove tRNA, residual proteins, EtBr and other host contaminants. These steps are not scaleable and therefore not suitable for use in large-scale processing. In contrast, preparative scale chromatography is a powerful purification tool that provides high resolution, operational ease and increased productivity for purifying DNA plasmid products. Two different modes of chromatography, reversed phase and anion exchange, show suitability in purifying DNA plasmid to the stringent levels required for human use. Separations based on reversed phase are governed by hydrophobic interactions while those for anion exchange are based on electrostatic interaction. These two orthogonal chromatography steps achieve separations between various forms of plasmid (supercoiled, open relaxed, linear and concatemers) and remove

- 8 -

host contaminants like LPS (endotoxin), RNA, DNA and residual proteins.

In the process of the present invention, harvested microbial cells are resuspended in modified STET buffer which is comprised of about 50 mM TRIS, about 50-100 mM EDTA, about 8% Sucrose, about 2% TRITON X-100, and optionally sub-microgram concentrations of lysozyme, at a pH in the range of 6.0-10.0. The concentration of lysozyme optionally used in the process of the present invention is substantially less than the concentration of lysozyme used in the procedures known in the art. It is readily apparent to those skilled in the art that modifications of this basic buffer formula can be made and are suitable for use in the present invention. Modifications to this basic buffer formula that do not substantially affect or alter the outcome of the present process are intended to be within the scope of the process of the present invention. The pH range may be adjusted according to the best results provided for the particular strain of bacteria being used. The preferred pH range is about 8.0-8.5. The suspension is then heated to about 70-100°C, with about 70-77°C preferred, in a flow-through heat exchanger. The lysate is centrifuged to pellet large cell debris, protein and most genomic DNA.

A prototype heat exchanger was built to demonstrate the feasibility of flow-through heat lysis of microbial cells containing plasmid. The particular heat exchanger consisted of a 10 ft. x 0.25 inch O.D. stainless steel tube shaped into a coil. The coil was completely immersed into a constant high temperature water bath. The hold-up volume of the coil was about 50 mL. Thermocouples and a thermometer were used to measure the inlet and exit temperatures, and the water bath temperature, respectively. The product stream was pumped into the heating coil using a Masterflex peristaltic pump with silicone tubing. Cell lysate exited the coil and was then centrifuged in a Beckman J-21 batch centrifuge for clarification. Figure 1 provides a schematic of this particular apparatus, however other types of heat exchanger construction are suitable for use in the present invention,

- 9 -

including but not limited to a shell and tube construction, which is preferable.

After centrifugation, the clarified lysate can optionally be treated with RNase, and the plasmid product can be filtered to further
5 remove small debris. A wide variety of filtration means are suitable for use in this process, including but not limited to filtration through a membrane having a small pore size. A preferred filtration method is filtration through a 0.45 micron filter.

To further remove contaminants from the DNA product,
10 the material can be diafiltered. Standard, commercially available diafiltration materials are suitable for use in this process, according to standard techniques known in the art. A preferred diafiltration method is diafiltration using an ultrafilter membrane having a molecular weight cutoff in the range of 30,000 to 500,000, depending on the plasmid size.
15 The DNA preparation described above is diafiltered using an ultrafiltration membrane (about 100,000 molecular weight cutoff) against column buffer prior to loading on the anion exchange column. Diafiltration prior to the anion exchange column is preferred, and it greatly increases the amount of lysate that can be loaded onto the
20 column.

A wide variety of commercially available anion exchange matrices are suitable for use in the present invention, including but not limited to those available from POROS Anion Exchange Resins, Qiagen, Toso Haas, Sterogene, Spherodex, Nucleopac, and Pharmacia. The
25 column (Poros II PI/M, 4.5 mm x 100) is initially equilibrated with 20 mM Bis/TRIS Propane at pH 7.5 and 0.7 M NaCl. The sample is loaded and washed with the same initial buffer. An elution gradient of 0.5 M to 0.85 M NaCl in about 25 column volumes is then applied and fractions are collected. Anion exchange chromatography is an ideal
30 first polishing step because it provides excellent clearance of RNA, genomic DNA and protein. Figure 5 (bold) shows a sample elution profile of filtered clarified cell lysate from the anion exchange column. Agarose gel analysis revealed that the second peak which appears after the flow-through is composed of the plasmid product. The earlier large

- 10 -

peak is due to RNA. This is confirmed by incubating the clarified cell lysate with ribonuclease prior to loading on the column, which showed that the large peak disappears and is replaced by several smaller more rapidly eluting peaks, due to the degradation products of ribonuclease digestion.

The anion exchange product fraction is loaded onto a reversed phase column. A wide variety of commercially available matrices are suitable for use in the present invention, including but not limited to those available from POROS, Polymer Labs, Toso Haas, Pharmacia, PQ Corp., Zorbax, BioSeptra resins, BioSeptra Hyper D resins, BioSeptra Q-Hyper D resins and Amicon. The matrices can also be polymer based or silica based. The reversed phase column (Poros R/H), is equilibrated with about 100 mM Ammonium Bicarbonate at pH 8.5. A gradient of 0-11% isopropanol is then used to elute bound material. The three forms of plasmids, forms I, II and III described above, can be separated by this method.

The eluted plasmid DNA can then be concentrated and/or diafiltered to reduce the volume or to change the buffer. For DNA intended for human use it may be useful to diafilter the DNA product into a pharmaceutically acceptable carrier or buffer solution. Pharmaceutically acceptable carriers or buffer solutions are known in the art and include those described in a variety of texts such as Remington's Pharmaceutical Sciences. Any method suitable for concentrating a DNA sample is suitable for use in the present invention. Such methods includes diafiltration, alcohol precipitation, lyophilization and the like, with diafiltration being preferred. Following diafiltration the final plasmid DNA product may then be sterilized. Any method of sterilization which does not affect the utility of the DNA product is suitable, such as sterilization by passage through a membrane having a sufficiently small pore size, for example 0.2 microns and smaller.

The following examples are provided to illustrate the process of the present invention without, however, limiting the same thereto.

- 11 -

EXAMPLE 1

Growth of Microbial Cells, Cell Lysis and Clarification

5 One liter of frozen *E. coli* cell slurry was used to make 8
liters of cell suspension in STET buffer (8% sucrose, 0.5% TRITON,
50 mM TRIS buffer, pH 8.5 and 50 mM EDTA). The absorbance of
the cell suspension at 600 nm was about O.D. 30. The suspension was
10 stirred continuously to ensure homogeneity. The viscosity of the cell
suspension was measured to be about 1.94 cp at room temperature
(24°C). The cell suspension was pumped through the heat exchanger at
81 mL/min which corresponded to a residence time of the cell solution
in the heat exchanger of about 35 seconds. The bath temperature was
15 maintained at 92°C. The inlet and outlet temperatures of the cell
solution were measured to be about 24°C and about 89°C (average),
respectively. About 1 liter of sample was run through the heat
exchanger. No visible clogging of the tube was observed although the
lysate was somewhat thicker than the starting material. The lysate was
cooled to room temperature and its viscosity was measured to be about
20 40 cp. The cell lysate was clarified by batch centrifugation at 9000
RPM for 50 minutes using the Beckman J-21. Analysis of the
supernatant confirmed effective cell lysis and product recovery. The
product yield produced by flow-through heat lysis was at least
comparable to that made by the Quigley & Holmes boiling method. The
25 latter method; however, must be carried out at the laboratory scale in
batch mode and is therefore unsuitable for large-scale (5 liters or
greater) processing. Since the heat exchanger process is flow-through,
there is no maximum limit to the volume of cell suspension that can be
processed. This process can therefore accomodate very large scale
30 fermentations of bacteria to produce large quantities of highly purified
plasmid DNA.

 The clarified lysate was then filtered through a membrane
having a pore size of 0.45 microns to remove finer debris. The filtrate
was then diafiltered using a membrane having a molecular weight cutoff
35 of about 100,000.

- 12 -

EXAMPLE 2

Control and Reproducibility of Cell Lysis with the Heat Exchanger

Adjusting the flowrate (i.e., residence time) at which the cell slurry is pumped through the heat exchanger permits tight control of the temperature of lysis, i.e., the outlet temperature. A cell slurry solution was prepared as described in Example 1 and pumped through the heat exchanger at flow rates ranging from 160 to 850 mL/min. The corresponding outlet temperatures ranged between 93°C and 65°C, respectively. Figure 2 illustrates the relationship between flow rate and temperature. The initial temperature of the cell slurry was 24°C and the bath temperature was kept constant at 96°C. In addition, a number of runs were performed where an outlet temperature of 80°C was targeted. Yields of 24 mg of circular DNA per L of clarified supernatant were consistently obtained demonstrating the reproducibility of the process.

EXAMPLE 3

Purification of Plasmid DNA

Microbial cells and lysates were prepared as described in Examples 1 and 2, and the following analyses were performed.

To illustrate that the addition of 100 mM EDTA vs 50 mM EDTA increased the percentage of supercoiled DNA, and to determine an acceptable range of outlet temperatures (i.e., lysis temperature) with respect to recovery of supercoiled DNA, the following analyses were performed. The supercoiled form of plasmid DNA is desirable since it is more stable than the relaxed circle form. One way that supercoiled DNA can be converted to open circle is by nicking with DNase. We found that the addition of 100 mM EDTA vs 50 mM in the STET buffer minimized the formation of open circle plasmid. Figure 3 shows comparative chromatograms of the total plasmid in the clarified supernatant with 50 mM EDTA vs 100 mM EDTA. The cell suspension was prepared as described in Example 1. The operating flow rate for

- 13 -

these runs was approximately 186 ml/min. The temperatures of the inlet, outlet and bath are 24°C, 92°C and 96°C respectively.

An acceptable range of lysis temperatures was determined by measuring the percentage of supercoiled plasmid generated for each run. Figure 4 illustrates the concentration of supercoiled plasmid as a function of exit temperature. An acceptable range of lysis temperatures is between 75°C and 92°C. At temperatures below 75°C, more relaxed circle plasmid was generated, most likely due to increased DNase activity. Above 93°C, the levels of supercoiled plasmid appear to diminish, possibly due to heat denaturation.

Following continuous heat lysis and centrifugation, 1 mL of clarified lysate was either incubated with 5 µg RNase for 2 hours, or was used untreated. The RNase treated and untreated samples were then loaded onto an anion exchange column (Poros Q/M 4.6 X 100) that had been previously equilibrated with a 50-50 mixture of solvents A and B [HPLC solvent A: 20 mM Tris/Bis Propane, pH 8.0; and solvent B: 1 M NaCl in 20 mM Tris/Bis Propane, pH 8.0]. The column was eluted using a gradient of 50% to 85% B run over 100 column volumes. Open circle plasmid elutes at approximately 68% B and supercoiled elutes at 72% B.

A comparison of the anion exchange column eluate from clarified lysate treated with RNase (thin line) and untreated (thick line) is shown in Figure 5. The peak at about 10 minutes is plasmid DNA, and is followed by a large peak in the untreated sample which is RNA. In the RNase treated sample the large RNA peak has been eliminated and a greater separation of the plasmid peak from contaminant peaks is produced.

As described above, diafiltration prior to anion exchange chromatography greatly increases the amount of lysate that can be loaded onto the column. This is demonstrated in Figure 6 which shows a comparison of clarified lysate which was diafiltered and clarified lysate which was not diafiltered prior to anion exchange chromatography. Samples were prepared as described above except that one sample was diafiltered before loading onto the anion exchange column,

- 14 -

and the other sample was not diafiltered. The column was run and eluted as described above. Figure 6 shows that the amount of contaminant material eluted from the column is vastly greater in the sample that was not diafiltered. The large amount of contaminating material which binds the anion exchange column matrix can overwhelm the maximum capacity of the column causing loss of DNA product because of the unavailability of the matrix to bind any more material. Therefore, diafiltration removes contaminants and allows more of the DNA product to bind the anion exchange matrix, and in turn allows a greater volume of clarified lysate to be loaded onto the column.

The plasmid DNA eluted from the anion exchange column was separated into the individual forms by reversed phase HPLC analysis. The separation of supercoiled plasmid (form 1) from nicked circle (form 2) is shown in Figure 7. The two forms were easily separated and allowed the isolation of individual forms of the plasmid.

EXAMPLE 4

Highly Purified Plasmid DNA From a Chromatography-based Process

A fermentation cell paste was resuspended in modified STET buffer and then thermally lysed in a batchwise manner. Alternatively a fermentation cell paste is resuspended in modified STET buffer and then thermally lysed in the flow-through process described above. The lysate was centrifuged as described above. Twenty ml of the supernatant were filtered as described above and loaded onto an anion exchange column (Poros Q/M 4.6x100) that was previously equilibrated with a 50-50 mixture of buffers A and B described above. A gradient of 50% to 85% B was run over 50 column volumes with a flow rate of 10ml/minute. Fractions of 2.5ml each were collected from the column. The supercoiled plasmid DNA eluted from the column at 72% B.

The anion exchange product was then loaded onto a reversed phase chromatography column (Poros R/H) which had been previously equilibrated with 100mM ammonium bicarbonate at pH 8.0,

- 15 -

and a gradient of 0% to 80% methanol was used to elute the bound material. The highly purified supercoiled plasmid DNA eluted at 22% methanol.

5 An agarose gel of the product fractions from each of the major steps of the purification process is shown in Figure 8. Based on the agarose gels and the colorimetric and HPLC assays described in Example 3, the final product, shown in Figure 9, is highly pure. The product consists of greater than 90% supercoiled and less than 10% open circle plasmid. RNA was below the limits of detection of the assay
10 used. Genomic DNA and protein contaminant levels were also below the limits of detection in the assays used. The overall supercoiled plasmid yield at the end of the process was approximately 60% of the supercoiled plasmid in the clarified lysate.

15

EXAMPLE 5

Multi-Gram Scale Purification of Plasmid DNA

4.5 L of frozen *E. coli* cell slurry was used to make 33.7 L of cell suspension in STET buffer (8% sucrose, 2% Triton, 50 mM Tris
20 buffer, 50 mM EDTA, pH 8.5) with 2500 units/ml of lysozyme. The absorbance of the suspension at 600 nm was about O.D. 30. The suspension was stirred at room temperature for 15 minutes to ensure proper mixing and then was incubated for 45 minutes with continuous stirring at 37°C. Following incubation, mixing was continued at room
25 temperature and the cell suspension was pumped through the heat exchanger at a flowrate of 500 ml/min. The batch temperature was maintained at 100°C and the inlet and outlet temperatures of the cell suspension were measured to be about 24°C and between 70-77°C, respectively. The cell lysate exiting the heat exchanger was collected in
30 Beckman centrifuge bottles (500 mls each) and the material was centrifuged immediately in Beckman J-21 centrifuges for 50 minutes at 9000 RPM. Following centrifugation, the supernatant was found to contain 4-5 times more plasmid product than in the case without lysozyme incubation. The supernatant product of the centrifugation was
35 immediately diafiltered against 3 volumes of TE buffer (25 mM Tris-

- 16 -

EDTA at pH 8.0) and then incubated with 20×10^5 units of *E. coli* RNase for 2-4 hours at room temperature. After completion of the incubation, the product solution was then diafiltered an additional 6 volumes with TE buffer using a 100 kD MWCO membrane and then filtered through a 0.45 micron filter to remove residual debris. The filtered lysate was diluted to 0.7 M NaCl with a 20 mM Bis/Tris Propane-NaCl buffer at pH 7.5, which prepares the diluted filtrate for loading onto the anion exchange column. The anion exchange column (3.6 L of POROS PI/M) was previously equilibrated with 20 mM Bis/Tris Propane and 0.7M NaCl. The filtered lysate was loaded to column capacity. In this case 5 grams of supercoiled plasmid was loaded onto the anion exchange column. After loading, the column was washed with 2-4 column volumes of 20 mM Bis/Tris Propane and 0.7 M NaCl. A 10 column volume gradient from 0.7 M NaCl to 2.0 M NaCl in 20 mM Bis/Tris Propane was performed to clear most of the *E. coli* protein. RNA and some endotoxin. The supercoiled plasmid fraction eluted between 1.4 M and 2.0 M NaCl. The supercoiled fraction from the anion exchange column, which contained 4 grams of supercoiled plasmid was then diluted 2-3 times with pyrogen free water, adjusted to 1.2% IPA and pH adjusted to 8.5 with 1 N NaOH. The diluted anion exchange supercoiled fraction was then loaded onto a 7 L reversed phase column (POROS R2/M) which had been previously equilibrated with 100 mM Ammonium Bicarbonate containing 1.2% IPA. In this case, 3.2 grams of supercoiled plasmid were loaded onto the reversed phase column and then the column was washed with 6-10 column volumes of 1.2% IPA in 100 mM Ammonium Bicarbonate. This extensive wash was performed to clear impurities. Next, a gradient of 1.2% IPA to 11.2% IPA in 5 column volumes was performed. The supercoiled plasmid fraction elutes at about 4% IPA. The supercoiled product fraction from the reversed phase column was then concentrated and diafiltered into normal saline using a 30 kD MWCO membrane. The final product bulk was filtered through a 0.22 micron filter. Table 1 provides a purification table describing clearance of impurities and yields across each of the major process steps. The overall product yield of the

- 17 -

process was more than 50% of the supercoiled plasmid in the clarified cell lysate as indicated by the anion exchange HPLC assay described in EXAMPLE 3. The purity of the product was very high with less than 1% *E. coli* RNA and protein, and less than 2.9% *E. coli* genomic DNA.

5

TABLE 1MULTI-GRAM PURIFICATION AND RECOVERY SUMMARY

10

STEP	Plasmid Product (mg)	% Step Yield	genomic DNA (mg/mg)	Protein (mg/mg)	RNA (mg/mg)	LAL Eu/mg
clarified lysate	6750	100	0.52	7.6	196	1.1x10 ⁷
concentration/RNase/ diafiltration/dead-end filtration	6500	93	0.50	1.6	2.21	3.4x10 ⁶
anion exchange	4000 of 5000	80	0.41	0.3	0.1	1.2x 10 ⁴
reversed phase	2300 of 3200	77	0.029	<0.01‡	<0.01‡	62
concentration diafiltration into final buffer	2110	100	0.029	<0.01‡	<0.01‡	2.8
final process yield		54				

‡ below detection limits of assay method

15

- 18 -

WHAT IS CLAIMED IS:

1. A process for large scale isolation and purification of plasmid DNA from large scale microbial cell fermentations,
- 5 comprising:
- a) harvesting microbial cells from a large scale fermentation;
 - b) adding to the harvested microbial cells a
10 sufficient amount of a lysis solution;
 - c) heating the microbial cells of step b) to a temperature between 70°C and 100°C in a flow-through heat exchanger to form a crude
15 lysate;
 - d) centrifuging the crude lysate;
 - e) filtering and diafiltering the supernatant of
20 step d) providing a filtrate;
 - f) contacting the filtrate of step e) with an anion exchange matrix;
 - g) eluting and collecting plasmid DNA from the
25 anion exchange matrix;
 - h) contacting the plasmid DNA from step g) with a reversed phase high performance liquid
30 chromatography matrix;
 - i) eluting and collecting the plasmid from the reversed phase high performance liquid chromatography matrix of step h);

- 19 -

j) optionally concentrating and/or diafiltering the product of step i) into a pharmaceutically acceptable carrier; and

k) optionally sterilizing the DNA product.

2. The process of Claim 1 wherein the lysis solution of step b) is modified STET buffer.

3. The process of Claim 1 wherein the heating of step c) is to a temperature between 70°C and 77°C.

15 4. The process of Claim 1 wherein the lysis solution of step b) contains a sub-microgram concentration of lysozyme.

5. The process of Claim 1 optionally including RNase treatment at any step following step a).

20 6. An isolated and purified plasmid DNA obtained by
the process of Claim 1.

7. The plasmid DNA of Claim 6 wherein said plasmid is suitable for administration to humans.

8. The plasmid DNA of Claim 6 wherein said plasmid is suitable for administration to non-human animals.

9. The plasmid DNA of Claim 6 wherein said plasmid is
30 a polynucleotide vaccine.

10. An isolated and purified plasmid DNA suitable for administration to humans.

- 20 -

11. The plasmid DNA of Claim 10 wherein said plasmid DNA is a polynucleotide vaccine.

1/9




FIG.1

2 / 9




FIG.2

3/9




FIG. 3

4/9




FIG.4

5/9




FIG. 5

6/9




FIG. 6

7/9




FIG. 7

8/9




FIG.8

9/9




FIG. 9

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/07083

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/10 C12P19/34

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BIOTECHNIQUES (1988), 6(9), 834, 837-8 CODEN: BTNQDQ; ISSN: 0736-6205, XP000600195 LE BRUN, J. J. ET AL: "A simple method for the preparation of the covalently closed circular form of plasmid DNA" see the whole document ---	6-8, 10
X	SAMBROOK ET AL.: "Molecular cloning" 1989, CSH LABORATORY PRESS, COLD SPRING HARBOR XP002011439	6-8
Y	see page 1.34 - page 1.35 ---	1, 2, 4, 5
	-/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

2 September 1996

Date of mailing of the international search report

06.09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gurdjian, D

INTERNATIONAL SEARCH REPORT

In' tional Application No
PCT/US 96/07083

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BIOTECHNIQUES (1992), 12(3), 430-4 CODEN: BTNQDQ;ISSN: 0736-6205, XP002012269 HINES, RONALD N. ET AL: "Large-scale purification of plasmid DNA by anion-exchange high - performance liquid chromatography"	6-10
Y	see abstract see page 433, right-hand column, paragraph 2	1,2,4,5
X	--- BIO-SCIENCES (1989), 8(1-2), 42-5 CODEN: BIOSER;ISSN: 0292-8418, XP000579530 COLOTE, S. ET AL: "Use of high-performance liquid chromatography in genetic engineering"	6-8
Y	see the whole document	1,2,4,5
X	--- WO,A,95 07995 (APPLIED IMMUNESCIENCES) 23 March 1995 see claims 55-28,33-43,	10,11
X	--- WO,A,91 06309 (UNIV VANDERBILT) 16 May 1991 see page 5, paragraph 2; claims 1-10	10
X	--- WO,A,93 24640 (UNIV CALIFORNIA ;DEBS ROBERT J (US); ZHU NING (US)) 9 December 1993 see claims 1,7,13	10
P,X	--- WO,A,96 02658 (MERCK & CO INC ;LEE ANN L (US); SAGAR SANGEETHA (US)) 1 February 1996 see the whole document -----	1-11

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/07083

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9507995	23-03-95	AU-B- 7717594 CA-A- 2171318 EP-A- 0719343 NO-A- 961010	03-04-95 23-03-95 03-07-96 07-05-96
WO-A-9106309	16-05-91	AU-B- 625013 AU-B- 6645690 CA-A- 2044593 EP-A- 0452457 JP-T- 4502772	25-06-92 31-05-91 04-05-91 23-10-91 21-05-92
WO-A-9324640	09-12-93	AU-B- 3467193 AU-B- 4407593 AU-B- 4528493 CA-A- 2126101 CA-A- 2134773 EP-A- 0625207 EP-A- 0646178 JP-T- 7507450 JP-T- 7502510 WO-A- 9312240 WO-A- 9325673	19-07-93 30-12-93 04-01-94 24-06-93 09-12-93 23-11-94 05-04-95 24-08-95 16-03-95 24-06-93 23-12-93
WO-A-9602658	01-02-96	AU-B- 3126295	16-02-96